arXiv:1303.3650v1 [math.AG] 15 Mar 2013

A Class of Incomplete Character Sums *

Lei Fu
Chern Institute of Mathematics and LPMC, Nankai University, Tianjin 300071, P. R. China
leifu@nankai.edu.cn

Daqing Wan
Department of Mathematics, University of California, Irvine, CA 92697, USA
dwan@math.uci.edu

Abstract

We define the transfer of lisse $\overline{\mathbb{Q}}_\ell$-sheaves of rank 1, and use it to study a class of incomplete character sums.

Key words: character sum, transfer, ℓ-adic cohomology.
Mathematics Subject Classification: 11L40, 11T23,

Introduction

Through out this paper, p is a fixed prime number, $\mathbb{F}_p$ is a finite field with p elements, and $\mathbb{F}$ is an algebraic closure of $\mathbb{F}_p$. For any power q of p, let $\mathbb{F}_q$ be the subfield of $\mathbb{F}$ with q elements. Let ℓ be a prime number distinct from p, let $\chi_1, \ldots, \chi_k : \mathbb{F}_{q^d}^* \to \overline{\mathbb{Q}}_\ell^*$ be a family of multiplicative characters and $\psi : \mathbb{F}_{q^d} \to \overline{\mathbb{Q}}_\ell^*$ an additive character on an extension field $\mathbb{F}_{q^d}$ of $\mathbb{F}_q$. We extend χ_i $(i = 1, \ldots, k)$ to $\mathbb{F}_{q^d}$ by setting $\chi_i(0) = 0$. Let $f_1(t_1, \ldots, t_n), \ldots, f_k(t_1, \ldots, t_n) \in \mathbb{F}_{q^d}[t_1, \ldots, t_n]$ be a family of polynomials with coefficients in $\mathbb{F}_{q^d}$, and let $f_{k+1}(t_1, \ldots, t_n) \in \mathbb{F}_{q^d}(t_1, \ldots, t_n)$ be a rational function. Motivated by a number of applications in [6], we are interested in estimating the following type of incomplete character sum

$$S' = \sum_{x_1,\ldots,x_n \in \mathbb{F}_q, f_{k+1}(x_1,\ldots,x_n) \neq \infty} \chi_1\big(f_1(x_1, \ldots, x_n)\big) \cdots \chi_k\big(f_k(x_1, \ldots, x_n)\big) \psi\big(f_{k+1}(x_1, \ldots, x_n)\big),$$

where the summation is over those $x_1, \ldots, x_n \in \mathbb{F}_q$ such that $f_{k+1}(x_1, \ldots, x_n)$ is defined. Note that for the classical complete character sum

$$S = \sum_{x_1,\ldots,x_n \in \mathbb{F}_{q^d}, f_{k+1}(x_1,\ldots,x_m) \neq \infty} \chi_1\big(f_1(x_1, \ldots, x_n)\big) \cdots \chi_k\big(f_k(x_1, \ldots, x_n)\big) \psi\big(f_{k+1}(x_1, \ldots, x_n)\big),$$

---

*The research of Lei Fu is supported by the NSFC (10525107).

the summation is over all $x_1, \ldots, x_n \in \mathbb{F}_{q^d}$ such that $f_{k+1}(x_1, \ldots, x_n)$ is defined. Under suitable hypothesis, one can give a sharp estimate of the following form for the complete sum

$$|S| \leq C(n, D)\sqrt{q}^{nd},$$

where $C(n, D)$ is a constant depending only on n and the degree D of the functions f_i's. We believe that there is also a sharp estimate for the incomplete sum S'. Namely, under suitable hypothesis, there should be an estimate of the form

$$|S'| \leq C(n, d, D)\sqrt{q}^{n},$$

where $C(n, d, D)$ is a constant depending only on n, d and the degree D. This is indeed true in the one variable case $n = 1$, see [6] and [4]. In the present paper, we use the transfer and ℓ-adic cohomology to treat the higher dimensional case.

Let us recall the general method of studying (complete) character sum using ℓ-adic cohomology theory. Let X_0 be a separated scheme of finite type over $\mathbb{F}_q$, let $X = X_0 \otimes_{\mathbb{F}_q} \mathbb{F}$ be the base change of X_0, let $\mathcal{L}_0$ be a lisse $\overline{\mathbb{Q}}_\ell$-sheaf on X_0, where ℓ is a prime different from the characteristic p. Let $\mathcal{L}$ be the inverse image of $\mathcal{L}_0$ in X. By definition, the L-function of $\mathcal{L}_0$ is defined to be

$$L(X_0, \mathcal{L}_0, t) = \prod_{x \in |X_0|} \frac{1}{\det(1 - F_x t^{\deg(x)}, \mathcal{L}_{0,\bar{x}})},$$

where $|X_0|$ is the set of Zariski closed points in X_0, and for any $x \in |X_0|$, $\deg(x)$ is the degree of the residue field $k(x)$ of $\mathcal{O}_{X,x}$ over the ground field $\mathbb{F}_q$, $\mathcal{L}_{0,\bar{x}}$ is the stalk of $\mathcal{L}_0$ at a geometric point over x, and $F_x \in \mathrm{Gal}(\overline{k(x)}/k(x))$ is the geometric Frobenius element, that is, the inverse of the Frobenius substitution. The family of complete character sums corresponding to $\mathcal{L}_0$ is the family of sums

$$S_m(X_0, \mathcal{L}_0) = \sum_{x \in X_0(\mathbb{F}_{q^m})} \mathrm{Tr}(F_x, \mathcal{L}_{0,\bar{x}}),$$

where $X_0(\mathbb{F}_{q^m}) = \mathrm{Hom}_{\mathrm{Spec}\,\mathbb{F}_q}(\mathrm{Spec}\,\mathbb{F}_{q^m}, X_0)$ is the set of $\mathbb{F}_{q^m}$-points in X_0. One can verify that

$$t\frac{d}{dt} \ln L(X_0, \mathcal{L}_0, t) = \sum_{m=1}^{\infty} S_m(X_0, \mathcal{L}_0) t^m.$$

So the L-function $L(X_0, \mathcal{L}_0, t)$ and the family of character sums $S_m(X_0, \mathcal{L}_0)$ $(m = 1, 2, \ldots)$ determine each other. By Grothendieck's trace formula ([1, Rapport 3.2]), we have

$$S_m(X_0, \mathcal{L}_0) = \sum_{i=0}^{2\dim(X)} (-1)^i \mathrm{Tr}\big(F^m, H_c^i(X, \mathcal{L})\big),$$

where $H_c^i(X, \mathcal{L})$ are the ℓ-adic cohomology groups with compact support and F is the geometric Frobenius correspondence. Deligne's theorem [2, Corollarie 3.3.4] gives us estimate for the Archimedean absolute values of the eigenvalues of F on $H_c^i(X, \mathcal{L})$. If we have some information about the Betti numbers $\dim H_c^i(X, \mathcal{L})$, then we can get estimate for $S_m(X_0, \mathcal{L}_0)$.

Let $\mathbb{F}_{q^d}$ be an extension of $\mathbb{F}_q$ of degree d, let $X_1 = X_0 \otimes_{\mathbb{F}_q} \mathbb{F}_{q^d}$ be the base change of X_0, and let $\mathcal{L}_1$ be a lisse $\overline{\mathbb{Q}}_\ell$-sheaf on X_1. For any $\mathbb{F}_q$-point $x : \operatorname{Spec} \mathbb{F}_q \to X_0$ of X_0, let $x_1 : \operatorname{Spec} \mathbb{F}_{q^d} \to X_1$ be its base change. It is a $\mathbb{F}_{q^d}$-point of X_1. In this paper, we are concerned with the following incomplete character sum

$$S_1'(X_1, \mathcal{L}_1) = \sum_{x \in X_0(\mathbb{F}_q)} \operatorname{Tr}(F_{x_1}, \mathcal{L}_{1,\bar{x}_1}),$$

where the summation is over those $\mathbb{F}_{q^d}$-points of X_1 arising from $\mathbb{F}_q$-points of X_0 by base change. We do not know how to estimate such incomplete sum in general. In the case where the sheaf $\mathcal{L}_1$ has rank 1, we construct the transfer $\operatorname{tran}(\mathcal{L}_1)$ of $\mathcal{L}_1$, which is a lisse $\overline{\mathbb{Q}}_\ell$-sheaf of rank 1 on X_0, so that $S_1'(X_1, \mathcal{L}_1)$ coincides with the complete character sum

$$S_1(X_0, \operatorname{tran}(\mathcal{L}_1)) = \sum_{x \in X_0(\mathbb{F}_q)} \operatorname{Tr}(F_x, (\operatorname{tran}(\mathcal{L}_1))_{\bar{x}}).$$

Theoretically, this reduces the study of our incomplete sum to the study of the complete sum for the new sheaf $\operatorname{tran}\mathcal{L}_1$. The problem is that it is hard to estimate the Betti numbers $\dim H_c^i(X, \operatorname{tran}(\mathcal{L}_1))$. In §4, we study the case where $\dim(X) = 1$. In this case, the Grothendieck-Ogg-Shararevich formula for Euler characteristic numbers of sheaves on curves gives us more information about the Betti numbers. We can thus get relatively complete results (Theorems 2.2-4).

Note that in [3, §3], using the so-called partial L-functions, we study a class of incomplete character sums including sums studied in this paper. The result in [3] applies to the case where $\mathcal{L}_1$ is of higher rank, but $\mathcal{L}_1$ is required to be able to descend down to a sheaf on X_0. In this paper, $\mathcal{L}_1$ may not be able to descend down, but it must be of rank 1.

1 Transfer

Let G be a pro-finite group and let H be an open subgroup of G of index d. Choose representatives $Hg_1, \ldots, Hg_d$ for the family of right cosets $H \backslash G$. For any $g \in G$, the set $\{Hg_1 g, \ldots, Hg_d g\}$ is also a family of representatives of right cosets. So we have

$$Hg_i g = Hg_{\tau(i)}$$

for a permutation $i \mapsto \tau(i)$ of $\{1, \ldots, d\}$. Define the *transfer* to be

$$\mathrm{tran} : G^{\mathrm{ab}} \to H^{\mathrm{ab}}, \quad g\overline{[G,G]} \to \prod_{i=1}^{d} g_i g g_{\tau(i)}^{-1} \overline{[H,H]},$$

where $G^{\mathrm{ab}} = G/\overline{[G,G]}$ and $H^{\mathrm{ab}} = H/\overline{[H,H]}$ are the largest abelian quotients of G and H (in the category of profinite groups), respectively. One can show that tran is independent of the choice of representatives of right cosets and is a well-defined homomorphism. The following proposition follows directly from the definition of transfer.

Proposition 1.1. *Let G be a pro-finite group and let H be an open subgroup of G of index d.*

(i) Then composite

$$G^{\mathrm{ab}} \overset{\mathrm{tran}}{\to} H^{\mathrm{ab}} \to G^{\mathrm{ab}}$$

coincides with the homomorphism $g\overline{[G,G]} \to g^d\overline{[G,G]}$, where the second arrow $H^{\mathrm{ab}} \to G^{\mathrm{ab}}$ in the composite is induced by the inclusion $H \hookrightarrow G$.

(ii) Suppose that H is normal in G. Let $\{Hg_1, \ldots, Hg_d\}$ be a family of representatives for $H\backslash G$. Then for any $h \in H$, we have

$$\mathrm{tran}(h\overline{[G,G]}) = \prod_{i=1}^{d} g_i h g_i^{-1} \overline{[H,H]}.$$

(iii) Suppose that H is normal in G and G/H is a cyclic group. For any $\sigma \in G$ such that σH is a generator of G/H, we have

$$\mathrm{tran}(\sigma\overline{[G,G]}) = \sigma^d \overline{[H,H]}.$$

Let X_0 be a scheme of finite type over $\mathbb{F}_q$, and let $X_1 = X_0 \otimes_{\mathbb{F}_q} \mathbb{F}_{q^d}$ be its base change to an extension field $\mathbb{F}_{q^d}$ of degree d over $\mathbb{F}_q$. Suppose X_0 is geometrically connected, that is, the base change $X = X_0 \otimes_{\mathbb{F}_q} \mathbb{F}$ is connected. Fix a geometric point for X. Take its images in X_0 and X_1 as the base points of X_0 and X_1, respectively, and let $\pi_1(X_0)$ and $\pi_1(X_1)$ be the etale fundamental groups with respect to these base points. Then $\pi_1(X_1)$ is a normal subgroup of $\pi_1(X_0)$. We have an isomorphism

$$\pi_1(X_0)/\pi_1(X_1) \overset{\cong}{\to} \mathrm{Gal}(\mathbb{F}_{q^d}/\mathbb{F}_q),$$

and $\mathrm{Gal}(\mathbb{F}_{q^d}/\mathbb{F}_q)$ is a cyclic group. Let $x : \mathrm{Spec}\,\mathbb{F}_{q^m} \to X_0$ be an $\mathbb{F}_{q^m}$-points in X_0. We can talk about the geometric Frobenius element F_x in $\pi_1(X_0)$ corresponding to x. It is defined up to conjugation in $\pi_1(X_0)$, and hence its image in $\pi_1(X_0)^{\mathrm{ab}}$ is well-defined. Sometimes we denote this image also by F_x

by abuse of notation. Consider the case where $m = 1$. Then the image of F_x in the quotient group $\pi_1(X_0)/\pi_1(X_1)$ is a generator. By Proposition 1.1 (iii), we have

$$\mathrm{tran}(F_x\overline{[\pi_1(X_0), \pi_1(X_0)]}) = F_x^d\overline{[\pi_1(X_1), \pi_1(X_1)]}.$$

Let $x_1 : \operatorname{Spec} \mathbb{F}_{q^d} \to X_1$ be the base change of x. One can verify that F_x^d and F_{x_1} define the same conjugacy class in $\pi_1(X_1)$. So we have $\mathrm{tran}(F_x\overline{[\pi_1(X_0), \pi_1(X_0)]}) = F_{x_1}\overline{[\pi_1(X_1), \pi_1(X_1)]}$, which we write as

$$\mathrm{tran}(F_x) = F_{x_1}$$

by abuse of notation.

Let $\mathcal{L}_1$ be a lisse $\overline{\mathbb{Q}}_\ell$-sheaf on X_1 of rank 1. It defines a one dimensional representation

$$\pi_1(X_1)^{\mathrm{ab}} \xrightarrow{\mathcal{L}_1} \overline{\mathbb{Q}}_\ell^*.$$

Taking its composite with the transfer $\mathrm{tran} : \pi_1(X_0)^{\mathrm{ab}} \to \pi_1(X_1)^{\mathrm{ab}}$, we get a one dimensional representation

$$\pi_1(X_0)^{\mathrm{ab}} \xrightarrow{\mathrm{tran}} \pi_1(X_1)^{\mathrm{ab}} \xrightarrow{\mathcal{L}_1} \overline{\mathbb{Q}}_\ell^*.$$

We define $\mathrm{tran}(\mathcal{L}_1)$ to be the lisse $\overline{\mathbb{Q}}_\ell$-sheaf on X_0 of rank 1 corresponding to the last representation. By the above discussion, if $x : \operatorname{Spec} \mathbb{F}_q \to X_0$ is an $\mathbb{F}_q$-point in X_0 and $x_1 : \operatorname{Spec} \mathbb{F}_{q^d} \to X_1$ is its base change, then we have

$$\mathrm{Tr}\big(F_x, (\mathrm{tran}(\mathcal{L}_1))_{\bar{x}}\big) = \mathrm{Tr}(F_{x_1}, \mathcal{L}_{1,\bar{x}_1}).$$

The following proposition follows from this equality, Proposition 1.1 (ii), and the definition of transfer.

Proposition 1.2. *Keep the notation above.*

(i) The incomplete character sum

$$S_1'(X_1, \mathcal{L}_1) = \sum_{x\in X_0(\mathbb{F}_q)} \mathrm{Tr}(F_{x_1}, \mathcal{L}_{1,\bar{x}_1})$$

coincides with the complete character sum

$$S_1(X_0, \mathrm{tran}(\mathcal{L}_1)) = \sum_{x\in X_0(\mathbb{F}_q)} \mathrm{Tr}\big(F_x, (\mathrm{tran}(\mathcal{L}_1))_{\bar{x}}\big).$$

(ii) Let $\rho_1 : \pi_1(X_1) \to \overline{\mathbb{Q}}_\ell^*$ be the character corresponding to the $\overline{\mathbb{Q}}_\ell$-sheaf $\mathcal{L}_1$ of rank 1, and let $\sigma \in \pi_1(X_0)$ be an element such that its image under the homomorphism $\pi_1(X_0) \to \mathrm{Gal}(\mathbb{F}_{q^d}/\mathbb{F}_q)$ is the Frobenius substitution. For each $0 \le i \le d-1$, let $\rho_1^{(i)} : \pi_1(X_1) \to \overline{\mathbb{Q}}_\ell^*$ be the character defined by

$$\rho_1^{(i)}(g) = \rho_1(\sigma^i g \sigma^{-i}),$$

and let $\pi : X_1 \to X_0$ be the projection. Then the character corresponding the sheaf $\pi^*(\mathrm{tran}(\mathcal{L}_1))$ on X_1 is $\prod_{i=0}^{d-1} \rho_1^{(i)}$.

(iii) Let $\mathcal{L}_1$ and $\mathcal{M}_1$ be two lisse $\overline{\mathbb{Q}}_\ell$-sheaves of rank 1 on X_1. Then we have

$$\mathrm{tran}(\mathcal{L}_1 \otimes \mathcal{M}_1) \cong \mathrm{tran}(\mathcal{L}_1) \otimes \mathrm{tran}(\mathcal{M}_1).$$

2 Transfer of Kummer type sheaves

The Kummer covering

$$[q-1] : \mathbb{G}_{m,\mathbb{F}_q} \to \mathbb{G}_{m,\mathbb{F}_q}, \quad x \mapsto x^{q-1}$$

on $\mathbb{G}_{m,\mathbb{F}_q} = \operatorname{Spec} \mathbb{F}_q[t, t^{-1}]$ defines a $\mathbb{F}_q^*$-torsor

$$1 \to \mathbb{F}_q^* \to \mathbb{G}_{m,\mathbb{F}_q} \overset{[q-1]}{\to} \mathbb{G}_{m,\mathbb{F}_q} \to 1.$$

Let $\chi : \mathbb{F}_q^* \to \overline{\mathbb{Q}}_\ell^*$ be a multiplicative character. Pushing-forward the above torsor by χ^{-1}, we get a lisse $\overline{\mathbb{Q}}_\ell$-sheaf $\mathcal{K}_\chi$ on $\mathbb{G}_{m,\mathbb{F}_q}$ of rank 1. We call $\mathcal{K}_\chi$ the *Kummer sheaf* associated to χ. For any $x \in \mathbb{G}_{m,\mathbb{F}_q}(\mathbb{F}_{q^m}) = \mathbb{F}_{q^m}^*$, we have

$$\mathrm{Tr}(F_x, \mathcal{K}_{\chi,\bar{x}}) = \chi(\mathrm{N}_{\mathbb{F}_{q^m}/\mathbb{F}_q}(x)),$$

where $\mathrm{N}_{\mathbb{F}_{q^m}/\mathbb{F}_q}$ denotes the norm for the field extension $\mathbb{F}_{q^m}/\mathbb{F}_q$.

Let $f \in \Gamma(X_0, \mathcal{O}_{X_0}^*)$ be a section of the subsheaf of units in the structure sheaf $\mathcal{O}_{X_0}$. Then the $\mathbb{F}_q$-algebra homomorphism

$$\mathbb{F}_q[t, t^{-1}] \to \Gamma(X_0, \mathcal{O}_{X_0}), \quad t \mapsto f$$

defines an $\mathbb{F}_q$-morphism $X_0 \to \mathbb{G}_{m,\mathbb{F}_q}$ which we still denote by f. Denote $f^*\mathcal{K}_\chi$ by $\mathcal{K}_{\chi,f}$. It is a lisse $\overline{\mathbb{Q}}_\ell$-sheaf of rank one on X_0. For any $x \in X_0(\mathbb{F}_{q^m})$, we have

$$\mathrm{Tr}(F_x, (\mathcal{K}_{\chi,f})_{\bar{x}}) = \chi\big(\mathrm{N}_{\mathbb{F}_{q^m}/\mathbb{F}_q}(f(x))\big).$$

For any $f_1, f_2 \in \Gamma(X_0, \mathcal{O}_{X_0}^*)$, we have

$$\mathcal{K}_{\chi,f_1 f_2} \cong \mathcal{K}_{\chi,f_1} \otimes \mathcal{K}_{\chi,f_2}.$$

Confer [1, Sommes trig. 1.4-8].

Suppose that X_0 is a normal scheme. Then $\mathcal{K}_{\chi,f}$ can be described as follows. Let K be the function field of X_0. Fix a separable closure of $\overline{K}$ of $K(X_0)$. Let the base point of X_0 be the canonical morphism $\operatorname{Spec}\overline{K} \to X_0$. Then we have a canonical isomorphism

$$\pi_1(X_0) \cong \varprojlim_{L} \operatorname{Gal}(L/K),$$

where L goes over those finite Galois extensions of K contained in $\overline{K}$ such that the normalization of X_0 in L is etale over X_0. (Actually it suffices to require that the normalization is unramified over X_0.) Let $\xi \in \overline{K}$ be a root of the polynomial $T^{q-1} - f$. Then any root of $T^{q-1} - f$ is of the form $a\xi$ for some $a \in \mathbb{F}_q^*$, $K[\xi]$ is a Galois extension, and the normalization of X_0 in $K[\xi]$ is etale over X_0. It follows that we have a canonical epimorphism

$$\pi_1(X_0) \to \operatorname{Gal}(K[\xi]/K).$$

On the other hand, we have a canonical monomorphism

$$\operatorname{Gal}(K[\xi]/K) \hookrightarrow \mathbb{F}_q^*, \quad g \mapsto \frac{g(\xi)}{\xi}.$$

The representation corresponding to $\mathcal{K}_{\chi,f}$ is the composite

$$\pi_1(X_0) \to \operatorname{Gal}(K[\xi]/K) \hookrightarrow \mathbb{F}_q^* \overset{\chi^{-1}}{\to} \overline{\mathbb{Q}}_\ell^*.$$

Let $\mathbb{F}_{q^d}$ be an extension of $\mathbb{F}_q$ of degree d, let $\chi : \mathbb{F}_{q^d}^* \to \overline{\mathbb{Q}}_\ell^*$ be a multiplicative character, let $f \in \mathbb{F}_{q^d}[t_1, \ldots, t_n]$, and let X_0 be an open subscheme of $\mathbb{A}^n_{\mathbb{F}_q}$ so that its inverse image in $\mathbb{A}^n_{\mathbb{F}_{q^d}}$ is contained in the complement of the hypersurface $f = 0$. Let X_1 be the base change of X_0 from $\mathbb{F}_q$ to $\mathbb{F}_{q^d}$, and let $\pi : X_1 \to X_0$ be the projection. Consider the sheaf $\mathcal{K}_{\chi,f}$ on X_1. Let us describe the character $\pi_1(X_1) \to \overline{\mathbb{Q}}_\ell^*$ corresponding to the sheaf $\pi^*(\operatorname{tran}(\mathcal{K}_{\chi,f}))$ on X_1. The function field of X_0 (resp. X_1) is the rational function field $K_0 = \mathbb{F}_q(t_1, \ldots, t_n)$ (resp. $K_1 = \mathbb{F}_{q^d}(t_1, \ldots, t_n)$). Let ξ be a root of the polynomial $T^{q^d-1} - f$ in a separable closure $\overline{K}_1$ of $K_1 = \mathbb{F}_{q^d}(t_1, \ldots, t_n)$. Then $\mathcal{K}_{\chi,f}$ corresponds to the character

$$\pi_1(X_1) \to \operatorname{Gal}(K_1[\xi]/K_1) \hookrightarrow \mathbb{F}_{q^d}^* \overset{\chi^{-1}}{\to} \overline{\mathbb{Q}}_\ell^*.$$

Denote this character by $\rho_1 : \pi_1(X_1) \to \overline{\mathbb{Q}}_\ell^*$. More explicitly, we have

$$\rho_1(g) = \chi^{-1}\left(\frac{g(\xi)}{\xi}\right)$$

for any $g \in \pi_1(X_1)$. Note that $\pi : X_1 \to X_0$ is a Galois etale covering space, and we have canonical isomorphisms

$$\operatorname{Aut}(X_1/X_0)^\circ \cong \operatorname{Gal}\Big(\mathbb{F}_{q^d}(t_1, \ldots, t_n)/\mathbb{F}_q(t_1, \ldots, t_n)\Big) \cong \operatorname{Gal}(\mathbb{F}_{q^d}/\mathbb{F}_q).$$

Choose an element $\sigma \in \pi_1(X_0)$ so that its image in $\mathrm{Gal}\Big(\mathbb{F}_{q^d}(t_1,\ldots,t_n)/\mathbb{F}_q(t_1,\ldots,t_n)\Big)$ is given by

$$\sigma(a) = a^q \text{ for any } a \in \mathbb{F}_{q^d}, \quad \sigma(t_i) = t_i \ (i = 1,\ldots,n).$$

Then the image of σ in $\pi_1(X_0)/\pi_1(X_1)$ is a generator. For any $0 \le i \le d-1$, let $\rho_1^{(i)} : \pi_1(X_1) \to \overline{\mathbb{Q}}_\ell^*$ be the character defined by

$$\rho_1^{(i)}(g) = \rho_1(\sigma^i g \sigma^{-i})$$

for any $g \in \pi_1(X_1)$. We have

$$\begin{aligned}\rho_1^{(i)}(g) &= \rho_1(\sigma^i g \sigma^{-i}) \\ &= \chi^{-1}\left(\frac{\sigma^i g \sigma^{-i}(\xi)}{\xi}\right) \\ &= \chi^{-1}\left(\sigma^i\left(\frac{g\sigma^{-i}(\xi)}{\sigma^{-i}(\xi)}\right)\right).\end{aligned}$$

Note that $\sigma^{-i}(\xi)$ and $g\sigma^{-i}(\xi)$ are roots of the polynomial $T^{q^d-1} - \sigma^{-i}(f)$, and hence $\frac{g\sigma^{-i}(\xi)}{\sigma^{-i}(\xi)}$ lies in $\mathbb{F}_{q^d}^*$. We thus have

$$\sigma^i\left(\frac{g\sigma^{-i}(\xi)}{\sigma^{-i}(\xi)}\right) = \left(\frac{g\sigma^{-i}(\xi)}{\sigma^{-i}(\xi)}\right)^{q^i}.$$

So we have

$$\rho_1^{(i)}(g) = (\chi^{q^i})^{-1}\left(\frac{g\sigma^{-i}(\xi)}{\sigma^{-i}(\xi)}\right)$$

for any $g \in \pi_1(X_1)$. This shows that $\rho_1^{(i)} : \pi_1(X_1) \to \overline{\mathbb{Q}}_\ell^*$ corresponds to the sheaf $\mathcal{K}_{\chi^{q^i},\sigma^{-i}(f)}$.We also have

$$\sigma^i\left(\frac{g\sigma^{-i}(\xi)}{\sigma^{-i}(\xi)}\right) = \left(\frac{g\sigma^{-i}(\xi)}{\sigma^{-i}(\xi)}\right)^{q^i} = \frac{g\Big(\big(\sigma^{-i}(\xi)\big)^{q^i}\Big)}{\big(\sigma^{-i}(\xi)\big)^{q^i}},$$

and hence

$$\rho_1^{(i)}(g) = \chi^{-1}\left(\frac{g\Big(\big(\sigma^{-i}(\xi)\big)^{q^i}\Big)}{\big(\sigma^{-i}(\xi)\big)^{q^i}}\right)$$

for any $g \in \pi_1(X_1)$. Note that $\big(\sigma^{-i}(\xi)\big)^{q^i}$ is a root of the polynomial $T^{q^d-1} - f(t_1^{q^i},\ldots,t_n^{q^i})$. This shows that $\rho_1^{(i)} : \pi_1(X_1) \to \overline{\mathbb{Q}}_\ell^*$ also corresponds to the sheaf $\mathcal{K}_{\chi,f(t_1^{q^i},\ldots,t_n^{q^i})}$. By Proposition 1.2 (ii), $\pi^*(\mathrm{tran}(\mathcal{K}_{\chi,f}))$ corresponds to the character $\prod_{i=0}^{d-1}\rho_1^{(i)}$. So we get the following.

Proposition 2.1. *Let* $\chi : \mathbb{F}_{q^d}^* \to \overline{\mathbb{Q}}_\ell^*$ *be a multiplicative character, let* $f \in \mathbb{F}_{q^d}[t_1,\ldots,t_n]$*, and let* X_0 *be an open subscheme of* $\mathbb{A}^n_{\mathbb{F}_q}$ *so that its inverse image in* $\mathbb{A}^n_{\mathbb{F}_{q^d}}$ *is contained in the complement of the* *hypersurface* $f = 0$. *Let* X_1 *be the base change of* X_0 *from* $\mathbb{F}_q$ *to* $\mathbb{F}_{q^d}$, *and let* $\pi : X_1 \to X_0$ *be the projection. Then on* X_1, *we have isomorphisms*

$$\pi^*(\mathrm{tran}(\mathcal{K}_{\chi,f})) \cong \bigotimes_{i=0}^{d-1} \mathcal{K}_{\chi^{q^i},\sigma^{-i}(f)} \cong \mathcal{K}_{\chi,\prod_{i=0}^{d-1} f(t_1^{q^i},\ldots,t_n^{q^i})},$$

where $\sigma^{-i}(f)$ *is the polynomial obtained from* f *by taking the* q^i*-th root for each coefficient of* f.

3 Transfer of Artin-Schreier type sheaves

The Artin-Schreier covering

$$\wp : \mathbb{A}^1_{\mathbb{F}_q} \to \mathbb{A}^1_{\mathbb{F}_q}, \quad x \mapsto x^q - x$$

defines an $\mathbb{F}_q$-torsor

$$0 \to \mathbb{F}_q \to \mathbb{A}^1_{\mathbb{F}_q} \xrightarrow{\wp} \mathbb{A}^1_{\mathbb{F}_q} \to 0.$$

Let $\psi : \mathbb{F}_q \to \overline{\mathbb{Q}}_\ell^*$ be a nontrivial additive character. Pushing-forward this torsor by ψ^{-1}, we get a lisse $\overline{\mathbb{Q}}_\ell$-sheaf $\mathcal{L}_\psi$ of rank 1 on $\mathbb{A}^1_{\mathbb{F}_q}$, which we call the *Artin-Schreier sheaf.* For any $x \in \mathbb{A}^1_{\mathbb{F}_q}(\mathbb{F}_{q^m}) = \mathbb{F}_{q^m}$, we have

$$\mathrm{Tr}(F_x, \mathcal{L}_{\psi,\bar{x}}) = \psi(\mathrm{Tr}_{\mathbb{F}_{q^m}/\mathbb{F}_q}(x)),$$

where $\mathrm{Tr}_{\mathbb{F}_{q^m}/\mathbb{F}_q}$ denotes the trace for the field extension $\mathbb{F}_{q^m}/\mathbb{F}_q$.

Let $f \in \Gamma(X_0, \mathcal{O}_{X_0})$ be a section of the structure sheaf $\mathcal{O}_{X_0}$. Then the $\mathbb{F}_q$-algebra homomorphism

$$\mathbb{F}_q[t] \to \Gamma(X_0, \mathcal{O}_{X_0}), \quad t \mapsto f$$

defines an $\mathbb{F}_q$-morphism $X_0 \to \mathbb{A}^1_{\mathbb{F}_q}$ which we still denote by f. Denote $f^*\mathcal{L}_\psi$ by $\mathcal{L}_{\psi,f}$. It is a lisse $\overline{\mathbb{Q}}_\ell$-sheaf of rank one on X_0. For any $x \in X_0(\mathbb{F}_{q^m})$, we have

$$\mathrm{Tr}(F_x, (\mathcal{L}_{\chi,f})_{\bar{x}}) = \psi\big(\mathrm{Tr}_{\mathbb{F}_{q^m}/\mathbb{F}_q}(f(x))\big).$$

For any $f_1, f_2 \in \Gamma(X_0, \mathcal{O}^*_{X_0})$, we have

$$\mathcal{L}_{\psi,f_1+f_2} \cong \mathcal{L}_{\psi,f_1} \otimes \mathcal{L}_{\psi,f_2}.$$

Confer [1, Sommes trig. 1.4-8].

Suppose that X_0 is a normal scheme. Then $\mathcal{L}_{\psi,f}$ can be described as follows. Let K be the function field of X_0. Fix a separable closure of $\overline{K}$ of $K(X_0)$, and let the base point of X_0 be the canonical morphism $\mathrm{Spec}\,\overline{K} \to X_0$. Let $\zeta \in \overline{K}$ be a root of the polynomial $T^q - T - f$. Then any root of $T^q - T - f$ is of the form $\zeta + a$ for some $a \in \mathbb{F}_q$, $K[\zeta]$ is a Galois extension and the normalization of X_0 in $K[\zeta]$ is etale over X_0. It follows that we have a canonical epimorphism

$$\pi_1(X_0) \to \mathrm{Gal}(K[\zeta]/K).$$

On the other hand, we have a canonical monomorphism

$$\mathrm{Gal}(K[\zeta]/K) \hookrightarrow \mathbb{F}_q, \quad g \mapsto g(\zeta) - \zeta.$$

The representation corresponding to $\mathcal{L}_{\psi,f}$ is the composite

$$\pi_1(X_0) \to \mathrm{Gal}(K[\zeta]/K) \hookrightarrow \mathbb{F}_q \overset{\psi^{-1}}{\to} \overline{\mathbb{Q}}_\ell^*.$$

Again let $\mathbb{F}_{q^d}$ be an extension of $\mathbb{F}_q$ of degree d, let $\psi : \mathbb{F}_{q^d} \to \overline{\mathbb{Q}}_\ell^*$ be an additive character, let $f \in \mathbb{F}_{q^d}(t_1, \ldots, t_n)$ be a rational function, and let X_0 be an open subscheme of $\mathbb{A}^n_{\mathbb{F}_q}$ so that its inverse image is contained in the complement of zero set of the denominator of f. Let X_1 be the base change of X_0 from $\mathbb{F}_q$ to $\mathbb{F}_{q^d}$, and let $\pi : X_1 \to X_0$ be the projection. Consider the sheaf $\mathcal{L}_{\psi,f}$ on X_1. Let us describe the character $\pi_1(X_1) \to \overline{\mathbb{Q}}_\ell^*$ corresponding to the sheaf $\pi^*(\mathrm{tran}(\mathcal{L}_{\psi,f}))$ on X_1. The function field of X_0 (resp. X_1) is the rational function field $K_0 = \mathbb{F}_q(t_1, \ldots, t_n)$ (resp. $K_1 = \mathbb{F}_{q^d}(t_1, \ldots, t_n)$). Let ζ be a root of the polynomial $T^{q^d} - T - f$ in a separable closure $\overline{K}_1$ of $K_1 = \mathbb{F}_{q^d}(t_1, \ldots, t_n)$. Then $\mathcal{L}_{\psi,f}$ corresponds to the character

$$\pi_1(X_1) \to \mathrm{Gal}(K_1[\zeta]/K_1) \hookrightarrow \mathbb{F}_{q^d} \overset{\psi^{-1}}{\to} \overline{\mathbb{Q}}_\ell^*.$$

Denote this character by $\rho_1 : \pi_1(X_1) \to \overline{\mathbb{Q}}_\ell^*$. More explicitly, we have

$$\rho_1(g) = \psi^{-1}\big(g(\zeta) - \zeta\big)$$

for any $g \in \pi_1(X_1)$. We have canonical isomorphisms

$$\mathrm{Aut}(X_1/X_0)^\circ \cong \mathrm{Gal}\bigg(\mathbb{F}_{q^d}(t_1, \ldots, t_n)/\mathbb{F}_q(t_1, \ldots, t_n)\bigg) \cong \mathrm{Gal}(\mathbb{F}_{q^d}/\mathbb{F}_q).$$

Again choose $\sigma \in \pi_1(X_0)$ so that its image in $\mathrm{Gal}\bigg(\mathbb{F}_{q^d}(t_1, \ldots, t_n)/\mathbb{F}_q(t_1, \ldots, t_n)\bigg)$ is given by

$$\sigma(a) = a^q \text{ for any } a \in \mathbb{F}_{q^d}, \quad \sigma(t_i) = t_i \ (i = 1, \ldots, n).$$

For any $0 \leq i \leq d - 1$, let $\rho_1^{(i)} : \pi_1(X_1) \to \overline{\mathbb{Q}}_\ell^*$ be the character defined by

$$\rho_1^{(i)}(g) = \rho_1(\sigma^i g \sigma^{-i})$$

for any $g \in \pi_1(X_1)$. We have

$$\begin{aligned} \rho_1^{(i)}(g) &= \rho_1(\sigma^i g \sigma^{-i}) \\ &= \psi^{-1}\big(\sigma^i g \sigma^{-i}(\zeta) - \zeta\big) \\ &= \psi^{-1}\sigma^i\big(g\sigma^{-i}(\zeta) - \sigma^{-i}(\zeta)\big). \end{aligned}$$

Note that $\sigma^{-i}(\zeta)$ and $g\sigma^{-i}(\zeta)$ are roots of the polynomial $T^{q^d} - T - \sigma^{-i}(f)$, and hence $g\sigma^{-i}(\zeta) - \sigma^{-i}(\zeta)$ lies in $\mathbb{F}_{q^d}$. We thus have

$$\sigma^i\big(g\sigma^{-i}(\zeta) - \sigma^{-i}(\zeta)\big) = \big(g\sigma^{-i}(\zeta) - \sigma^{-i}(\zeta)\big)^{q^i}.$$

So we have

$$\rho_1^{(i)}(g) = \psi^{-1}\left(\big(g\sigma^{-i}(\zeta) - \sigma^{-i}(\zeta)\big)^{q^i}\right)$$

for any $g \in \pi_1(X_1)$. This shows that $\rho_1^{(i)} : \pi_1(X_1) \to \overline{\mathbb{Q}}_\ell^*$ corresponds to the sheaf $\mathcal{L}_{\psi\circ\sigma^i, \sigma^{-i}(f)}$. By Proposition 1.2 (ii), $\pi^*(\mathrm{tran}(\mathcal{L}_{\psi,f}))$ corresponds to the character $\prod_{i=0}^{d-1} \rho_1^{(i)}$. So we get the following.

Proposition 3.1. *Let $\psi : \mathbb{F}_{q^d} \to \overline{\mathbb{Q}}_\ell^*$ be an additive character, let $f \in \mathbb{F}_{q^d}(t_1, \ldots, t_n)$ be a rational function, and let X_0 be an open subscheme of $\mathbb{A}^n_{\mathbb{F}_q}$ so that its inverse image in $\mathbb{A}^n_{\mathbb{F}_{q^d}}$ is contained in the complement of zero set of the denominator of f. Let X_1 be the base change of X_0 from $\mathbb{F}_q$ to $\mathbb{F}_{q^d}$, and let $\pi : X_1 \to X_0$ be the projection. Then on X_1, we have an isomorphism*

$$\pi^*(\mathrm{tran}(\mathcal{L}_{\psi,f})) \cong \bigotimes_{i=0}^{d-1} \mathcal{L}_{\psi\circ\sigma^i, \sigma^{-i}(f)},$$

where $\sigma^{-i}(f)$ is the rational function obtained from f by taking the q^i-th root for each coefficient of the numerator and the denominator of f, and $\psi \circ \sigma^i$ is the additive character $a \mapsto \psi(a^{q^i})$ for any $a \in \mathbb{F}_{q^d}$.

For any rational function $f \in \mathbb{F}_{q^d}(t_1, \ldots, t_n)$, define

$$\mathrm{Tr}_{\mathbb{F}_{d^d}/\mathbb{F}_q}(f) = \sum_{i=0}^{d-1} \sigma^i(f).$$

Write $f = \frac{\sum_{i_1,\ldots,i_n} a_{i_1\ldots i_n} t_1^{i_1}\cdots t_n^{i_n}}{\sum_{i_1,\ldots,i_n} b_{i_1\ldots i_n} t_1^{i_1}\cdots t_n^{i_n}}$. Then we have

$$\mathrm{Tr}_{\mathbb{F}_{d^d}/\mathbb{F}_q}(f) = \sum_{i=0}^{d-1} \frac{\sum_{i_1,\ldots,i_n} a_{i_1\ldots i_n}^{q^i} t_1^{i_1}\cdots t_n^{i_n}}{\sum_{i_1,\ldots,i_n} b_{i_1\ldots i_n}^{q^i} t_1^{i_1}\cdots t_n^{i_n}}.$$

4 Character Sums on Curves

In this section, X_0 is a smooth geometrically connected curve over $\mathbb{F}_q$ of genus g, $\overline{X}_0$ is the smooth compactification of X_0, X and $\overline{X}$ are the base changes of X_0 and $\overline{X}_0$ from $\mathbb{F}_q$ to $\mathbb{F}$, respectively. Let us recall the general method of studying character sums on X_0 using ℓ-adic cohomology theory. Let $\mathcal{L}_0$ be a lisse $\overline{\mathbb{Q}}_\ell$-sheaf on X_0 of rank 1. Suppose $\mathcal{L}_0$ is *geometrically nontrivial*, that is, the inverse image $\mathcal{L}$ on X is not isomorphic to the constant sheaf $\overline{\mathbb{Q}}_\ell$. Then we have

$$H^0_c(X,\mathcal{L})=0,\quad H^2_c(X,\mathcal{L})=0.$$

By the Grothendieck-Ogg-Shafarevich formula ([5, X 7.1]), we have

$$\begin{aligned}\dim H^1_c(X,\mathcal{L}) &= -\sum_{i=0}^{2}(-1)^i\dim H^i_c(X,\mathcal{L})\\ &= \big(2g-2+\#(\overline{X}-X)\big)+\sum_{x\in\overline{X}-X}\mathrm{sw}_x(\mathcal{L}),\end{aligned}$$

where sw_x denotes the Swan conductor at x for any Zariski closed point x in $\overline{X}$. Suppose $\mathcal{L}$ is punctually pure of weight 0. (Confer [2, Définition 1.2.2]). Then by [2, Corollarie 3.3.4], all the eigenvalues of F on $H^1_c(X,\mathcal{L})$ have Archimedean absolute value $\leq q^{\frac{1}{2}}$. By the Grothendieck trace formula [1, Rapport 3.2], we have

$$\begin{aligned}S_m(X_0,\mathcal{L}_0) &= \sum_{x\in X_0(\mathbb{F}_{q^m})}\mathrm{Tr}(F_x,\mathcal{L}_{0,\bar{x}})\\ &= \sum_{i=0}^{2}(-1)^i\mathrm{Tr}\big(F^m,H^i_c(X,\mathcal{L})\big)\\ &= -\mathrm{Tr}\big(F^m,H^1_c(X,\mathcal{L})\big).\end{aligned}$$

So we have

$$|S_m(X_0,\mathcal{L}_0)|\leq\left(\big(2g-2+\#(\overline{X}-X)\big)+\sum_{x\in\overline{X}-X}\mathrm{sw}_x(\mathcal{L})\right)q^{\frac{m}{2}}.$$

We thus get the following.

Proposition 4.1. *Let X_0 be a smooth geometrically connected curve over $\mathbb{F}_q$ of genus g, $\overline{X}_0$ its smooth compactification, and $\mathcal{L}_0$ a lisse $\overline{\mathbb{Q}}_\ell$-sheaf on X_0 of rank 1. Suppose $\mathcal{L}_0$ is geometrically nontrivial,* *and is punctually pure of weight* 0*. Then we have*

$$\begin{aligned} |S_m(X_0,\mathcal{L}_0)| &= \left|\sum_{x\in X_0(\mathbb{F}_{q^m})}\mathrm{Tr}(F_x,\mathcal{L}_{0,\bar{x}})\right| \\ &\leq \left(\big(2g-2+\#(\overline{X}-X)\big)+\sum_{x\in\overline{X}-X}\mathrm{sw}_x(\mathcal{L})\right)q^{\frac{m}{2}}. \end{aligned}$$

Theorem 4.2. *Let* X_0 *be a smooth geometrically connected curve over* $\mathbb{F}_q$ *of genus* g*,* $\overline{X}_0$ *its smooth compactification,* $\mathbb{F}_{q^d}$ *an extension of* $\mathbb{F}_q$ *of degree* d*,* $X_1=X_0\otimes_{\mathbb{F}_q}\mathbb{F}_{q^d}$*, and* $\mathcal{L}_1$ *a lisse* $\overline{\mathbb{Q}}_\ell$*-sheaf on* X_1 *of rank* 1*. Suppose* $\mathrm{tran}(\mathcal{L}_1)$ *is geometrically nontrivial, and suppose* $\mathcal{L}_1$ *is punctually pure of weight* 0*. Let*

$$S_1'(X_1,\mathcal{L}_1)=\sum_{x\in X_0(\mathbb{F}_q)}\mathrm{Tr}(F_{x_1},\mathcal{L}_{1,\bar{x}_1}),$$

where $x_1:\mathrm{Spec}\,\mathbb{F}_{q^d}\to X_1$ *is the base change of* $x:\mathrm{Spec}\,\mathbb{F}_q\to X_0$ *for any* $x\in X_0(\mathbb{F}_q)$*. Then we have*

$$|S_1'(X_1,\mathcal{L}_1)|\leq\left(\big(2g-2+\#(\overline{X}-X)\big)+d\sum_{x\in\overline{X}-X}\mathrm{sw}_x(\mathcal{L})\right)\sqrt{q}.$$

Proof. By Proposition 1.2 (i) and Grothendieck's trace formula ([1, Rapport 3.2]), we have

$$\begin{aligned} S_1'(X_1,\mathcal{L}_1) &= \sum_{x\in X_0(\mathbb{F}_q)}\mathrm{Tr}\big(F_x,(\mathrm{tran}(\mathcal{L}_1))_{\bar{x}}\big) \\ &= \sum_{i=0}^{2}(-1)^i\mathrm{Tr}\big(F,H_c^i(X,\mathrm{tran}(\mathcal{L}_1))\big). \end{aligned}$$

Since $\mathrm{tran}(\mathcal{L}_1)$ is geometrically nontrivial, we have $H_c^i(X,\mathrm{tran}(\mathcal{L}_1))=0$ for $i\neq 1$. So we have

$$S_1'(X_1,\mathcal{L}_1)=-\mathrm{Tr}\big(F,H_c^1(X,\mathrm{tran}(\mathcal{L}_1))\big).$$

By [2, Corollarie 3.3.4], all eigenvalues of F on $H_c^1(X,\mathrm{tran}(\mathcal{L}_1))$ have Archimedean absolute value $\leq\sqrt{q}$. So we have

$$|S_1'(X_1,\mathcal{L}_1)|\leq\big(\dim H_c^1(X,\mathrm{tran}(\mathcal{L}_1))\big)\sqrt{q}.$$

On the other hand, by the Grothendieck-Ogg-Shafarevich formula ([5, X 7.1]), we have

$$\begin{aligned} \dim H_c^1(X,\mathrm{tran}(\mathcal{L}_1)) &= -\sum_{i=0}^{2}(-1)^i\dim H_c^i(X,\mathrm{tran}(\mathcal{L}_1)) \\ &= \big(2g-2+\#(\overline{X}-X)\big)+\sum_{x\in\overline{X}-X}\mathrm{sw}_x(\mathrm{tran}(\mathcal{L}_1)). \end{aligned}$$

To prove our theorem, it suffices to show that

$$\sum_{x\in\overline{X}-X}\mathrm{sw}_x(\mathrm{tran}(\mathcal{L}_1))\leq d\sum_{x\in\overline{X}-X}\mathrm{sw}_x(\mathcal{L}_1).$$

Choose $\sigma \in \pi_1(X_0)$ so that its image under the canonical epimorphism $\pi_1(X_0) \to \mathrm{Gal}(\mathbb{F}_{q^d}/\mathbb{F}_q)$ is the Frobenius substitution. Let $\rho_1 : \pi_1(X_1) \to \overline{\mathbb{Q}}_\ell^*$ be the character corresponding to $\mathcal{L}_1$ and let $\rho_1^{(i)} : \pi_1(X_1) \to \overline{\mathbb{Q}}_\ell^*$ $(0 \leq i \leq d-1)$ be the characters defined by

$$\rho_1^{(i)}(g) = \rho_1(\sigma^i g \sigma^{-i})$$

for any $g \in \pi_1(X_0)$. Then by Proposition 1.2, $\pi^*(\mathrm{tran}(\mathcal{L}_1))$ corresponds to the character $\prod_{i=0}^{d-1} \rho_1^{(i)}$, where $\pi : X_1 \to X_0$ is the projection. Note that σ induce an automorphism of X over X_0, and an automorphism of $\overline{X}$ over $\overline{X}_0$. For any $x \in \overline{X} - X$ and any $0 \leq i \leq d-1$, we have

$$\mathrm{sw}_x(\rho_1^{(i)}) = \mathrm{sw}_{\sigma^i(x)}(\rho_1).$$

So we have

$$\begin{aligned}
\mathrm{sw}_x(\mathrm{tran}(\mathcal{L}_1)) &= \mathrm{sw}_x\left(\prod_{i=0}^{d-1} \rho_1^{(i)}\right) \\
&\leq \max\{\mathrm{sw}_x(\rho_1^{(i)}) | 0 \leq i \leq d-1\} \\
&= \max\{\mathrm{sw}_{\sigma^i(x)}(\rho_1) | 0 \leq i \leq d-1\} \\
&= \max\{\mathrm{sw}_{\sigma^i(x)}(\mathcal{L}_1) | 0 \leq i \leq d-1\} \\
&\leq \sum_{i=0}^{d-1} \mathrm{sw}_{\sigma^i(x)}(\mathcal{L}_1).
\end{aligned}$$

Hence

$$\begin{aligned}
\sum_{x \in \overline{X} - X} \mathrm{sw}_x(\mathrm{tran}(\mathcal{L}_1)) &\leq \sum_{x \in \overline{X} - X} \sum_{i=0}^{d-1} \mathrm{sw}_{\sigma^i(x)}(\mathcal{L}_1) \\
&= d \sum_{x \in \overline{X} - X} \mathrm{sw}_x(\mathcal{L}_1).
\end{aligned}$$

□

Remark 4.3. In the above proof, suppose there exists closed points $\infty_1, \ldots, \infty_m \in \overline{X} - X$ such that $\sigma(\infty_j) = \infty_j$ $(i = 1, \ldots, m)$. Then we have

$$\begin{aligned}
\mathrm{sw}_{\infty_j}(\mathrm{tran}(\mathcal{L}_1)) &= \mathrm{sw}_{\infty_j}\left(\prod_{i=0}^{d-1} \rho_1^{(i)}\right) \\
&\leq \max\{\mathrm{sw}_{\infty_j}(\rho_1^{(i)}) | 0 \leq i \leq d-1\} \\
&= \max\{\mathrm{sw}_{\sigma^i(\infty_j)}(\rho_1) | 0 \leq i \leq d-1\} \\
&= \mathrm{sw}_{\infty_j}(\mathcal{L}_1).
\end{aligned}$$

The estimate in Theorem 4.2 can be improved as

$$|S_1'(X_1,\mathcal{L}_1)|\leq\left(\big(2g-2+\#(\overline{X}-X)\big)+d\sum_{x\in\overline{X}-X,\ x\neq\infty_1,\ldots,\infty_m}\mathrm{sw}_x(\mathcal{L})+\sum_{j=1}^m\mathrm{sw}_{\infty_j}(\mathcal{L}_1)\right)\sqrt{q}.$$

For any polynomial $f(t)$ in $\mathbb{F}_{q^d}[t]$, define the highest square free divisor of $f(t)$ to be the product of all distinct irreducible factors of $f(t)$. Let σ be the automorphism of $\mathbb{F}_{q^d}(t)$ defined by $\sigma(t)=t$ and $\sigma(a)=a^q$ for any $a\in\mathbb{F}_{q^d}$.

Theorem 4.4. *Let $f_1(t),\ldots,f_k(t)$ be pairwise relatively prime polynomials in $\mathbb{F}_{q^d}[t]$ whose highest square free divisors have degrees $d_1,\ldots,d_k$, respectively, and let $\chi_1,\ldots,\chi_k:\mathbb{F}_{q^d}^*\to\mathbb{C}^*$ be multiplicative characters for $\mathbb{F}_{q^d}$. Extend χ_j $(j=1,\ldots,k)$ to $\mathbb{F}_{q^d}$ by setting $\chi_j(0)=0$. Suppose that for some $1\leq j_0\leq k$, the polynomial $\prod_{i=0}^{d-1}f_{j_0}(t_1^{q^i},\ldots,t_n^{q^i})$ is not of the form $h(t)^{\mathrm{ord}(\chi_{j_0})}$ in $\mathbb{F}[t]$, where $\mathrm{ord}(\chi_{j_0})$ is the smallest integer d such that $\chi_{j_0}^d=1$. Then we have*

$$\left|\sum_{a\in\mathbb{F}_q}\chi_1(f_1(a))\cdots\chi_k(f_k(a))\right|\leq\left(d\sum_{j=1}^n d_j-1\right)\sqrt{q}.$$

Proof. Let X_0 be the complement in $\mathbb{A}^1_{\mathbb{F}_q}$ of the closed subscheme defined by the vanishing of the polynomial $\prod_{i=0}^{d-1}\prod_{j=1}^k\sigma^{-i}(f_j)$. Let $\mathcal{L}_1$ be the $\overline{\mathbb{Q}}_\ell$-sheaf

$$\mathcal{L}_1=\bigotimes_{j=1}^k\mathcal{K}_{\chi_j,f_j}$$

on $X_1=X_0\otimes_{\mathbb{F}_q}\mathbb{F}_{q^d}$. Then we have

$$\sum_{a\in\mathbb{F}_q}\chi_1(f_1(a))\cdots\chi_k(f_k(a))=\sum_{x\in X_0(\mathbb{F}_q)}\mathrm{Tr}(F_{x_1},\mathcal{L}_{1,\bar{x}_1}).$$

Note that $\mathcal{L}_1$ is tamely ramified everywhere on $\overline{X}_1=\mathbb{P}^1_{\mathbb{F}_{q^d}}$, and hence

$$\mathrm{sw}_x(\mathcal{L}_1)=0$$

for all $x\in\overline{X}-X=\mathbb{P}^1_{\mathbb{F}}-X_0\otimes_{\mathbb{F}_q}\mathbb{F}$. Note that the set $\overline{X}-X$ consists of ∞ and the roots of $\prod_{i=0}^{d-1}\prod_{j=1}^k\sigma^{-i}(f_j)$. There are at most $d\sum_{j=1}^n d_j$ such roots. The genus of X_0 is 0. Our estimate follows directly from the estimate in Theorem 4.2, provided that we can prove $\mathrm{tran}(\mathcal{L}_1)$ is geometrically nontrivial. Indeed, by Proposition 2.1, we have $\pi^*(\mathrm{tran}(\mathcal{K}_{\chi_{j_0},f_{j_0}}))\cong\mathcal{K}_{\chi_{j_0},\prod_{i=0}^{d-1}f_{j_0}(t_1^{q^i},\ldots,t_n^{q^i})}$. Since $\prod_{i=0}^{d-1}f_{j_0}(t_1^{q^i},\ldots,t_n^{q^i})$ is not of the form $h(t)^{\mathrm{ord}(\chi_{j_0})}$ in $\mathbb{F}[t]$, the sheaf $\mathcal{K}_{\chi_{j_0},\prod_{i=0}^{d-1}f_{j_0}(t_1^{q^i},\ldots,t_n^{q^i})}$ is geometrically nontrivial, and hence $\mathrm{tran}(\mathcal{K}_{\chi_{j_0},f_{j_0}})$ is geometrically nontrivial. Since f_{j_0} is relatively prime to f_j for any $j\neq j_0$, this implies that $\mathrm{tran}(\mathcal{L}_1)$ is geometrically nontrivial. $\square$

For any rational function $f = \frac{\sum_i a_i t^i}{\sum_i b_i t^i} \in \mathbb{F}_{q^d}(t)$, we define

$$\mathrm{Tr}_{\mathbb{F}_{d^d}/\mathbb{F}_q}(f) = \sum_{j=0}^{d-1} \frac{\sum_i a_i^{q^j} t^i}{\sum_i b_i^{q^j} t^i}.$$

Theorem 4.5. *Let $f_j(t)$ $(1 \leq j \leq k)$ be polynomials in $\mathbb{F}_{q^d}[t]$, let $f_{k+1}(t)$ be a rational function in $\mathbb{F}_{q^d}(t)$, let D_1 be the degree of the highest square free divisor of $\prod_{i=1}^k f_i(t)$, let $D_2 = 0$ if $\deg(f_{k+1}) \leq 0$ and $D_2 = \deg(f_{k+1})$ if $\deg(f_{k+1}) > 0$, let D_3 be the degree of the denominator of $f_{k+1}(t)$, and let D_4 be the degree of the highest square free divisor of the denominator of $f_{k+1}(t)$ which is relatively prime to $\prod_{j=1}^k f_j(t)$. Let χ_j $(1 \leq j \leq k)$ be multiplicative characters of $\mathbb{F}_{q^d}$, and let ψ_p be a nontrivial additive character of $\mathbb{F}_p$. Extend χ_i to $\mathbb{F}_{q^d}$ by setting $\chi_i(0) = 0$. Suppose $\mathrm{Tr}_{\mathbb{F}_{q^d}/\mathbb{F}_q}(f_{n+1}(t))$ is not of the form $r(t)^p - r(t) + c$ in $\mathbb{F}(t)$. Then we have the estimate*

$$\left| \sum_{a \in \mathbb{F}_q,\ f_{n+1}(a) \neq \infty} \chi_1(f_1(a)) \cdots \chi_n(f_n(a)) \psi_p \left(\mathrm{Tr}_{\mathbb{F}_{q^d}/\mathbb{F}_p}(f_{n+1}(a)) \right) \right| \leq (d(D_1 + D_3 + D_4) + D_2 - 1)\sqrt{q}.$$

Proof. Let u_{k+1} be the denominator of f_{k+1}, and let X_0 be the complement in $\mathbb{A}^1_{\mathbb{F}_q}$ of the closed subscheme defined by the vanishing of the polynomial $\prod_{i=0}^{d-1} \left(\sigma^{-i}(u_{k+1}) \prod_{j=1}^k \sigma^{-i}(f_j) \right)$. Let $\mathcal{L}_1$ be the $\overline{\mathbb{Q}}_\ell$-sheaf

$$\mathcal{L}_1 = \left(\bigotimes_{j=1}^k \mathcal{K}_{\chi_j, f_j} \right) \bigotimes \mathcal{L}_{\psi, f_{k+1}}$$

on $X_1 = X_0 \otimes_{\mathbb{F}_q} \mathbb{F}_{q^d}$, where $\psi = \psi_p \circ \mathrm{Tr}_{\mathbb{F}_{q^d}/\mathbb{F}_p}$. Then we have

$$\sum_{a \in \mathbb{F}_q} \chi_1(f_1(a)) \cdots \chi_k(f_k(a)) \psi_p \left(\mathrm{Tr}_{\mathbb{F}_{q^d}/\mathbb{F}_p}(f_{n+1}(a)) \right) = \sum_{x \in X_0(\mathbb{F}_q)} \mathrm{Tr}(F_{x_1}, \mathcal{L}_{1,\bar{x}_1}).$$

Note that $\bigotimes_{j=1}^k \mathcal{K}_{\chi_j, f_j}$ is tamely ramified everywhere on $\overline{X}_1 = \mathbb{P}^1_{\mathbb{F}_{q^d}}$, and has no effect on the Swan conductor of $\mathcal{L}_1$. So for any $x \in \overline{X} - X = \mathbb{P}^1_{\mathbb{F}} - X_0 \otimes_{\mathbb{F}_q} \mathbb{F}$, we have

$$\mathrm{sw}_x(\mathcal{L}_1) = \mathrm{sw}_x(\mathcal{L}_{\psi, f_{k+1}}).$$

Let v_x be the valuation of the field $\mathbb{F}(t)$ corresponding to the point $x \in \overline{X} - X$. If x is not a pole of f_{k+1}, then $\mathrm{sw}_x(\mathcal{L}_{\psi, f_{k+1}}) = 0$. Otherwise, we have

$$\mathrm{sw}_x(\mathcal{L}_{\psi, f_{k+1}}) \leq -v_x(f_{k+1}).$$

It follows that

$$\mathrm{sw}_\infty(\mathcal{L}) \leq D_2, \qquad \sum_{x \in \overline{X} - X,\ x \neq \infty} \mathrm{sw}_x(\mathcal{L}) \leq D_3.$$

The set $\overline{X} - X$ consists of ∞, roots of $\sigma^{-i}(f_j)$ $(0 \leq i \leq d-1,\ 1 \leq j \leq k)$, and roots of the denominators of $\sigma^{-i}(f_{k+1})$ $(0 \leq i \leq d-1)$. It follows that

$$\#(\overline{X} - X) \leq 1 + d(D_1 + D_4).$$

The genus of X_0 is 0. Our estimate follows directly from the estimate in Theorem 4.2 and Remark 4.3, provided that we can prove $\mathrm{tran}(\mathcal{L}_1)$ is geometrically nontrivial. Let $\pi : X_1 \to X_0$ be the projection. It suffices to prove $\pi^*(\mathrm{tran}(\mathcal{L}_1))$ is geometrically nontrivial. By Propositions 1.2 (iii), 2.1 and 3.1, we have

$$\pi^*(\mathrm{tran}(\mathcal{L}_1)) \cong \left(\bigotimes_{j=1}^{k} \bigotimes_{i=0}^{d-1} \mathcal{K}_{\chi_j^{q^i}, \prod_{j=1}^{k} \sigma^{-i}(f_j)} \right) \bigotimes \mathcal{L}_{\psi, \mathrm{Tr}_{\mathbb{F}_{q^d}/\mathbb{F}_q}(f_{k+1})}.$$

Since $\mathrm{Tr}_{\mathbb{F}_{q^d}/\mathbb{F}_q}(f_{k+1})$ is not of the form $r(t)^p - r(t) + c$ in $\mathbb{F}(t)$, the sheaf $\mathcal{L}_{\psi, \mathrm{Tr}_{\mathbb{F}_{q^d}/\mathbb{F}_q}(f_{k+1})}$ is geometrically nontrivial, and it has wild ramification at poles of $\mathrm{Tr}_{\mathbb{F}_{q^d}/\mathbb{F}_q}(f_{k+1})$, whereas $\bigotimes_{j=1}^{k} \bigotimes_{i=0}^{d-1} \bigotimes_{i=0}^{d-1} \mathcal{K}_{\chi_j^{q^i}, \prod_{j=1}^{d} \sigma^{-i}(f_k)}$ is at worst tamely ramified. So $\left(\bigotimes_{j=1}^{k} \bigotimes_{i=0}^{d-1} \bigotimes_{i=0}^{d-1} \mathcal{K}_{\chi_j^{q^i}, \prod_{j=1}^{k} \sigma^{-i}(f_j)} \right) \bigotimes \mathcal{L}_{\psi, \mathrm{Tr}_{\mathbb{F}_{q^d}/\mathbb{F}_q}(f_{k+1})}$ is geometrically nontrivial. $\square$

References

[1] P. Deligne, *Cohomologie Étale* (SGA $4\frac{1}{2}$), Lecture Notes in Math. 569, Springer-Verlag (1977).

[2] P. Deligne, La conjecture de Weil II, *Publ. Math. IHES* 52 (1980), 137-252.

[3] L. Fu and D. Wan, Moment L-functions, partial L-functions, and partial Exponential sums, *Math. Ann.* 328 (2004), 193-228.

[4] L. Fu and D. Wan, L-functions of character sums over finite fields, preprint.

[5] A. Grothendieck, *Cohomologie ℓ-adique et fonctions L* (SGA 5), Lecture Notes in Math. 589, Springer-Verlag (1977).

[6] D. Wan, Generators and irreducible polynomials over finite fields, Math. Comp., 66(1997), 1195-1212.